



## Original Article



## Comparison of Outcomes of Kidney Transplantation in Diabetic and Non-Diabetic Patients

Manzoor Ahmad Naeem<sup>1</sup>, Alia Naseer<sup>2\*</sup>, Sidra Naseem<sup>2</sup>, Mohammad Asad Bilal<sup>2</sup>, Usra Naeem<sup>3</sup>, Ali Raza<sup>2</sup> and Iqra Rehman<sup>2</sup>

<sup>1</sup>Department of Health Professional Technology, Faculty of Allied Health Sciences, Superior University, Lahore, Pakistan

<sup>2</sup>Department of Health Professional Technologies, Faculty of Allied Health Sciences, The University of Lahore, Lahore, Pakistan

<sup>3</sup>Department of Emerging Allied Health Technologies, Faculty of Allied Health Sciences, Superior University, Lahore, Pakistan

## ARTICLE INFO

## Keywords:

Kidney Transplantation, Diabetes Mellitus, Non-Diabetic Patients, End-Stage Renal Transplantation

## How to Cite:

Naeem, M. A., Naseer, A., Naseem, S., Bilal, M. A., Naeem, U., Raza, A., & Rehman, I. (2025). Comparison of Outcomes of Kidney Transplantation in Diabetic and Non-Diabetic Patients: Kidney Transplantation in Diabetic and Non-Diabetic Patients. *Pakistan BioMedical Journal*, 8(2), 21-24. <https://doi.org/10.54393/pbmj.v8i2.1185>

## \*Corresponding Author:

Alia Naseer

Department of Health Professional Technologies, Faculty of Allied Health Sciences, The University of Lahore, Lahore, Pakistan  
annanaseer@gmail.com

Received date: 27<sup>th</sup> December, 2024

Acceptance date: 15<sup>th</sup> February, 2025

Published date: 28<sup>th</sup> February, 2025

## ABSTRACT

When opposed to long-lasting dialysis, kidney transplantation offers patients with end-stage renal illness a higher chance of mortality and a better standard of life. It is still up for debate whether kidney transplantation is more beneficial for diabetic individuals in contrast to those lacking the disease. **Objective:** To compare outcomes of kidney transplantation in diabetic and non-diabetic patients. **Methods:** Data from 100 patients was collected from Sheikh Zayed Hospital Lahore, Pakistan. The following variables were assessed to compare the consequences of transplants in patients with and without diabetes: per graft fluid collection, superficial infection of the wounded area, cut-off hernia, shallow injury dehiscence, cellulitis, seroma, fascial breaking down, deep wound contamination, hypertension, and mortality and recuperation. Fisher's exact test was applied to test statistically significant variances in outcomes and  $p$ -values  $\leq 0.05$  was considered significant. **Results:** A statistically significant variance was observed in per graft fluid collection among diabetic and non-diabetic kidney transplant patients i.e., 68% vs 32% respectively with  $p$ -values  $\leq 0.05$ . A significant difference was observed in the occurrence of incisional hernias between the two groups, with a high rate in non-diabetic patients. Facial dehiscence also showed a statistically significant difference, with a higher incidence in non-diabetic patients ( $p=0.010$ ). **Conclusions:** It was concluded that diabetes significantly impacts certain post-transplant outcomes, such as fluid collection. In contrast, other outcomes like incisional hernia and facial dehiscence are more prevalent in non-diabetic patients who had kidney transplants.

## INTRODUCTION

When contrasted with long-lasting dialysis, kidney transplantation offers those suffering from end-stage renal disease (ESRD) a better standard of life and a higher chance of survival, perhaps saving their lives [1, 2]. However, there are several inconsistencies in earlier research, such as the fact that adherence to glycemic objectives was not taken into account and that diabetes therapy administered over time changed [3]. Because of the tiny sample size, our estimations are not as precise as possible. The regression models did not account for the incidence of medicated acute rejection or the Kidney Donor Profile Index. However, the success of a kidney transplant can be influenced by various factors, including

the recipient's underlying health conditions [4]. Diabetes mellitus (DM), a leading cause of ESRD, is particularly significant due to its pervasive effects on multiple organ systems, including the cardiovascular system, immune function, and wound healing [5, 6]. All but one study had a larger proportion of male respondents than female ones [7]. From Saudi Arabia to Australia, studies were carried out all over the world, although most came from China ( $n=6$ ) and Poland ( $n=3$ ). Diabetes was the condition most often evaluated result ( $n=14$ ) [8]. According to estimates, the prevalence of diabetes in all age categories was 2.8% worldwide in 2000 and is expected to increase to 4.4% by 2030 [9]. According to a 2006 research by the WHO



Collaborating Center for Diabetes and the Diabetic Association of Pakistan, the incidence of Type II diabetes was 2.5% in rural women and 6.9% in men, whereas it was 3.5% in women and 6.0% in men living in cities [10]. According to these standards, the incidence of diabetes worldwide was 8.8% (95% CI 7.2–11.3%) in 2017 (standardized for the 20–79 age range), and by 2045, it is predicted to rise to 9.9% (95% CI 7.5–12.7%). The incidence of diabetes has steadily increased worldwide [11]. The effectiveness of kidney transplantation in patients with diabetes as opposed to patients without diabetes is still debatable [12–14]. While some studies found that diabetic patients had poor graft survival and mortality, others found no appreciable differences between patients with and without diabetes [13, 14].

This study aims to compare the outcomes of kidney transplantation in diabetic (DM) and non-diabetic patients (NDM).

## METHODS

It was a retrospective study and data were collected from Sheikh Zayed Hospital Lahore after approval from the ethical review committee. The study duration was from Sep 2024 to Dec 2024. Ethical Approval was given by the ethical review board of The University of Lahore. (REC-UOL-/401/08/24). A sample size of 100 with a 95% confidence level and 5% margin of error was calculated using Rao software. The sample selection technique involved was non-probability purposive sampling from patients who had kidney transplants between 20–50 years of age. Both male and female patients were included. ABO conflicting or human leukocyte antigen-sensitized kidney transplantation and immunocompromised patients were excluded. After getting informed consent from patients a total of 100 patients were enrolled in this study. To compare outcomes of transplant in diabetic and non-diabetic patients' variables that were measured include: 1) per graft fluid collection, 2) superficial wound infection, 3) incision hernia, 4) superficial wound dehiscence, 5) cellulitis, seroma, 6) fascial dehiscence, 7) deep wound infection, 8) hypertension, 9) death and recovery. SPSS version 23.0 was used to analyze the data. The distinction between the final results of kidney transplant recipients with DM and those with NDM was described using frequency and percentage. To look for statistically significant variances in results, Fisher's exact test was applied and a p-value < 0.05 seemed significant.

## RESULTS

Out of 100 patients that had kidney transplantation 48% were in the age group of 41–50 years of age. Seventy-nine percent of patients were female. A statistically significant variance was investigated in per graft fluid collection among diabetic and non-diabetic kidney transplant patients i.e., 68% vs 32% respectively with p-value ≤ 0.05

(Table 1).

**Table 1:** Demographic Analysis of Diabetic Kidney Transplant Patients

Variables	%	p-value
<b>Gender</b>		
Female	79%	--
Male	21%	
<b>Age</b>		
41–50 Years	48%	--
<b>Graft Fluid Collection</b>		
Diabetic Kidney Transplant Patients	68%	≤0.05
Non-Diabetic Kidney Transplant Patients	32%	

A significant variance was noticed in the occurrence of incisional hernias between groups, with an elevated occurrence incidence in non-diabetic patients. Facial dehiscence also gives a statistically significant difference, with an elevated occurrence in non-diabetic patients (p=0.010). Six diabetic kidney transplant patients died compared to 3 non-diabetic patients (Table 2).

**Table 2:** Outcome of Kidney Transplant in DM and NDM Patients

Outcomes	DM	NDM	p-value
Superficial Dehiscence	52	43	0.530
Per Graft Fluid Collection	68	32	0.003
Superficial Wound Infection	61	39	0.086
Incisional Hernia	28	72	0.0003
Cellulitis	43	57	0.271
Seroma	39	61	0.086
Facial Dehiscence	34	66	0.010
Deep Wound Infection	59	41	0.177
Hypertension	51	49	0.903
Death	6	3	0.683
Recovery	44	47	0.898

## DISCUSSION

The outcomes of kidney transplant patients often vary based on underlying health conditions, with diabetes being a critical factor that can influence post-operative recovery and complications [15]. This research sought to understand the post-transplant outcomes between NDM and DM patients. Our findings showed considerable differences in some outcomes, emphasizing the unique challenges of diabetic patients after kidney transplantation. The frequency of per graft fluid collection was considerably higher in the DM group than in the NDM group (p=0.003). This implies that diabetic patients might be at increased risk of having fluid-related complications after transplant, based on diabetes's impact on wound healing and vascular integrity [16]. The incidence of incisional hernias was notably distinguished between the two groups, with the NDM group showing a higher rate. This result is surprising, considering a greater risk might be expected in diabetic patients based on more compromised tissue quality. It may represent variation in surgical

methods, post-operative care, or overall health of the patients [17]. For superficial dehiscence, superficial wound infection, cellulitis, seroma, deep wound infection, HTN, death, and recovery for other outcomes, there were no differences between DM and NDM groups. This implies that although diabetes mellitus affects some aspects of post-transplantation recovery, its effects may not be felt across all aspects [18, 19]. The comparable hypertension and overall recovery rates between the two groups show that both DM and NDM patients can have similar long-duration outcomes under proper management [20]. The results of this study highlight the need for individualized post-operative care in diabetic patients who undergo kidney transplantation. The substantially increased rates of fluid accumulation in the DM group imply that these patients may require more stringent observation and possibly more intensive fluid and wound management.

## CONCLUSIONS

It was concluded that diabetes significantly impacts certain post-transplant outcomes, such as fluid collection, in kidney transplant patients. In contrast, other outcomes like incisional hernia and facial dehiscence are more prevalent in non-diabetic patients. These findings highlight the need for nuanced post-operative care strategies that address the specific risks associated with both diabetic and non-diabetic patients.

## Authors Contribution

Conceptualization: MAN

Methodology: AN, SN, MAB

Formal analysis: AN, UN

Writing review and editing: SN, AR, IR

All authors have read and agreed to the published version of the manuscript.

## Conflicts of Interest

All the authors declare no conflict of interest.

## Source of Funding

The author received no financial support for the research, authorship and/or publication of this article.

## REFERENCES

- [1] Palleti SK, Avula S, Dewan S. Kidney: A Review on End Stage Renal Disease. *Dialysis and Transplant. International Clinical and Medical Case Reports Journal*.2023 Apr;2(10):doi:10.5281/zenodo.7791286.
- [2] Delautre A, Hannedouche T, Couchoud C, Guiserix J, Cerasuolo D, Chantrel F et al. Diabetic Kidney Disease Versus Non-Diabetic Kidney Disease in Type 2 Diabetic Patients On Dialysis: An Observational Cohort. *Endocrinology, Diabetes and Metabolism*. 2022 Jul;5(4): e00281. doi: 10.1002/edm2.281.
- [3] García-Padilla P, Dávila-Rúales V, Hurtado DC, Vargas DC, Muñoz OM, Jurado MA. A Comparative Study on Graft and Overall Survival Rates Between Diabetic and Nondiabetic Kidney Transplant Patients Through Survival Analysis. *Canadian Journal of Kidney Health and Disease*.2023 Sep; 10: 20543581231199011.doi: 10.1177/20543581231199011.
- [4] Wang JH, Skeans MA, Israni AK. Current Status of Kidney Transplant Outcomes:Dying to Survive. *Advances in Chronic Kidney Disease*.2016 Sep; 23(5): 281-6. doi: 10.1053/j.ackd.2016.07.001.
- [5] Mahbub T, Chowdhury MN, Jahan F, Arefeen S, Rahman M, Sarker MS. Diabetes is the Leading Cause of ESRD in Hemodialysis Patients.*Journal of Medicine*.2013; 14(1): 62.doi:10.3329/jom.v14i1.14595.
- [6] Daryabor G, Atashzar MR, Kabelitz D, Meri S, Kalantar K. The Effects of Type 2 Diabetes Mellitus On Organ Metabolism and the Immune System.*Frontiers in Immunology*.2020Jul;11:1582.doi:10.3389/fimmu.2020.01582.
- [7] Sayilar El, Ersoy A, Ersoy C, Oruc A, Ayar Y, Sigirli D. The Effect of Calcineurin Inhibitors On Anthropometric Measurements in Kidney Transplant Recipients. *BioMed Central Nephrology*.2022 Nov; 23(1): 375. doi: 10.1186/s12882-022-03004-1.
- [8] Kajdas AA, Kleibert M, Normann AK, Krasuski K, Linde DS, Szostak-Węgierek D. Immunosuppressive Therapy and Nutritional Diseases of Patients After Kidney Transplantation: A Systematic Review. *BioMed Central Nephrology*.2025 Jan; 26(1): 33. doi: 10.1186/s12882-025-03964-0.
- [9] Wild S, Roglic G, Green A, Sicree R, King H. Global Prevalence of Diabetes: Estimates for the Year 2000 and Projections for 2030. *Diabetes Care*.2004 May; 27(5): 1047-53. doi: 10.2337/diacare.27.5.1047.
- [10] Shera AS, Jawad F, Maqsood A. Prevalence of Diabetes in Pakistan. *Diabetes Research and Clinical Practice*.2007 May;76(2):219-22.doi:10.1016/j.diabres.2006.08.011.
- [11] Standl E, Khunti K, Hansen TB, Schnell O. The global epidemics of diabetes in the 21st century: Current situation and perspectives. *European Journal of Preventive Cardiology*.2019 Dec; 26(2\_suppl): 7-14. doi: 10.1177/2047487319881021.
- [12] Revanur VK, Jardine AG, Kingsmore DB, Jaques BC, Hamilton DH, Jindal RM. Influence of Diabetes Mellitus On Patient and Graft Survival in Recipients of Kidney Transplantation. *Clinical Transplantation*. 2001 Apr; 15(2): 89-94. doi:10.1034/j.1399-0012.2001.150202.x.
- [13] Jassal SV, Krahn MD, Naglie G, Zaltzman JS, Roscoe JM, Cole EH et al. Kidney Transplantation in the Elderly: A Decision Analysis. *Journal of the American*

- Society of Nephrology. 2003 Jan; 14(1): 187-96. doi: 10.1097/01.ASN.0000042166.70351.57.
- [14] Boucek P, Saudek F, Pokorna E, Vitko S, Adamec M, Koznarova R *et al.* Kidney Transplantation in Type 2 Diabetic Patients: A Comparison with Matched Non-Diabetic Subjects. *Nephrology Dialysis Transplantation*. 2002 Sep; 17(9): 1678-83. doi: 10.1093/ndt/17.9.1678.
- [15] Baek CH, Kim H, Baek SD, Jang M, Kim W, Yang WS *et al.* Outcomes of Living Donor Kidney Transplantation in Diabetic Patients: Age and Sex Matched Comparison with Non-Diabetic Patients. *The Korean Journal of Internal Medicine*. 2017 Aug; 33(2): 356. doi: 10.3904/kjim.2016.067.
- [16] Schiel R, Heinrich S, Steiner T, Ott U, Stein G. Long-Term Prognosis of Patients After Kidney Transplantation: A Comparison of Those with or Without Diabetes Mellitus. *Nephrology Dialysis Transplantation*. 2005 Mar; 20(3): 611-7. doi: 10.1093/ndt/gfh657.
- [17] Ojo AO, Morales JM, González-Molina M, Steffick DE, Luan FL, Merion RM *et al.* Comparison of the Long-Term Outcomes of Kidney Transplantation: USA Versus Spain. *Nephrology Dialysis Transplantation*. 2013 Jan; 28(1): 213-20. doi: 10.1093/ndt/gfs287.
- [18] Rischen-Vos J, Van der Woude FJ, Tegzess AM, Zwinderman AH, Gooszen HC, Van Den Akker PJ *et al.* Increased Morbidity and Mortality in Patients with Diabetes Mellitus After Kidney Transplantation as Compared with Non-Diabetic Patients. *Nephrology Dialysis Transplantation*. 1992 Jan; 7(5): 433-7. doi: 10.1093/oxfordjournals.ndt.a092162.
- [19] Rafi IK, Aktaruzzaman M, Rahman FS, Amin MR. Hypertension May Be Driving Force to Enhance Kidney Disorder. *International Journal of Research*. 2024 May; 11(5): 108-115. doi: 10.5281/zenodo.11180907
- [20] Comai G, Corradetti V, Bini C, Tondolo F, Hu L, Valente S *et al.* Histological Findings of Diabetic Kidneys Transplanted in Non-Diabetic Recipients: A Case Series. *International Urology and Nephrology*. 2023 Oct; 55(10): 2611-9. doi: 10.1007/s11255-023-03552-x.