



OPEN ACCESS

Original Article

Isolation and Identification of *Escherichia coli* and *Klebsiella* species using Chromogenic Agar from Drinking Water of District Thar, Sindh, Pakistan

Jawaid Mian Larik¹, Shagufta Jabeen^{1,2*}, Beenish Khanzada³, Naveen Qadeer¹, Humaira Naz⁴, Bisma Adil¹ and Shaista Naz⁵

¹Institute of Microbiology, University of Sindh, Jamshoro, Pakistan

²Institute of Bioscience, Universiti Putra Malaysia, Malaysia

³Institute of Biochemistry, University of Sindh, Jamshoro, Pakistan

⁴Centre for Pure and Applied Geology, University of Sindh, Jamshoro, Pakistan

⁵Department of Geography, University of Sindh, Jamshoro, Pakistan

ARTICLE INFO

Keywords:

Coliform, Pathogen, Filtration, Chromogenic Agar, Infection, Contamination, Drinking Water

How to Cite:

Larik, J. M., Jabeen, S., Khanzada, B., Qadeer, N., Naz, H., Adil, B., & Naz, S. (2025). Isolation and Identification of *Escherichia coli* and *Klebsiella* Species using Chromogenic Agar from Drinking Water of District Thar, Sindh, Pakistan: Identification of *Escherichia coli* and *Klebsiella* Species from drinking water. *Pakistan BioMedical Journal*, 8(12), 29-33. <https://doi.org/10.54393/pbmj.v8i12.1330>

***Corresponding Author:**

Shagufta Jabeen

Institute of Bioscience, Universiti Putra Malaysia, Malaysia

shagufta.jabeen@usindh.edu.pk

Received Date: 11th November, 2025

Revised Date: 25th December, 2025

Acceptance Date: 29th December, 2025

Published Date: 31st December, 2025

ABSTRACT

Most of the studies do not conduct bacteriological analysis of the Thar region drinking water, except for transportation and travel problems. **Objectives:** To identify the bacteriological profile of water at various regions of the Thar Desert, Sindh, Pakistan, concerning the presence of the indicator Coliform, i.e. *E. coli* and *Klebsiella* spp., on Rapid Media Chromogenic agar.

Methods: A total of 70 drinking water samples were collected in 5 talukas of District Tharparkar, namely, Islamkot, Nangrpar, Kalohi, Dahri, and Diplo between December 2021 and May 2021. Water samples were tested right after the collection, with the help of the membrane filtration method. MacConkey agar and Chromogenic agar and plates were inoculated with filters and incubated aerobically in 35-37°C, respectively, and kept for 24 hours. Bacterial identification tests were performed on the colonies on MacConkey agar plates and *E. coli* and *Klebsiella* spp. produced turquoise-blue colony and Mergenda color colony on chromogenic agar, respectively.

Results: The findings of the bacteriological test revealed that the water samples are not potable, and 47 (67%) samples contained bacteria. There were 28 (59%) coliform and 19 (41%) non-coliforms. *Escherichia coli* (18; 40%) was most common, followed by *Klebsiella* spp. (12; 27%), *Pseudomonas* spp. (9; 20%), *Salmonella* spp. and *Proteus* spp. were the least common (3; 6%) each. **Conclusions:** Chromogenic agar is a fast and convenient way of presumptive identification of bacterial contaminants in drinking water. *E. coli* and *Klebsiella* spp. were the largest bacterial communities in drinking water specimens of the Thar region.

INTRODUCTION

Tharparkar is the largest region of the desert region of Sindh, located in Pakistan. Tharparkar is found in 69°3'35 E and 71° 7' 47 E longitudes and 24° 9' 35 N and 25° 43' 6 N latitudes [1]. The only significant drinking water source is groundwater, which is a significant source of domestic, livestock, and agricultural usage. These groundwater sources in Southern Sindh, such as Tharparkar, contain a significant number of coliform bacteria, which implies that the groundwater poses a health hazard to the locals [2].

The polluted drinking water is the cause of numerous water-borne illnesses, most commonly dysentery, vomiting, and gastroenteritis [3]. Polluted water has also been reported in various parts of Pakistan that harbor coliform bacteria and other pathogenic microorganisms such as *Escherichia coli*, *Vibrio cholera*, *Salmonella* spp., and *Pseudomonas* spp; that cause gastroenteritis, cholera, typhoid fever, diarrhea, and other enteric diseases in children and adults who consume it [4-6]. The

contamination bacteria (Total coliforms and *Escherichia coli*) of various cities of Sindh, such as Nagarparkar, have been reported, signifying the intrusion of sewage water into the drinking water system [7-9]. Therefore, various procedures involving agar and liquid media employing various techniques such as the membrane-filtration method, multiple tube or most probable number (MPN) methods are well elaborated to identify the water contamination to isolate indicator microorganism in water [10, 11]. The traditional procedures are tedious and require more time to detect bacterial contamination [12, 13]. Nevertheless, the use of agar with the aid of chromogenic agar makes it quicker, and even the process of identifying indicator microorganisms in drinking water is easier. Safe drinking water is important to the health of the people, particularly in arid areas such as Tharparkar, where the primary source of water is through groundwater [14]. The scarcity of literature in the field of bacteriological quality in this area demonstrates that the new surveillance with the help of fast techniques is necessary. This paper sought to determine the quality of drinking water in the District of Tharparkar by identifying bacterial contamination and coliform bacteria, which are indicators of the presence of pathogens, through rapid chromogenic agar media. This study aimed to present the evidence of the possible health hazards of the local water and to be able to inform the strategies of enhancing water safety.

METHODS

A cross-sectional descriptive study of coliform contamination and antibiotic resistance in drinking water from five talukas of District Tharparkar. The samples included a total of five talukas of District Tharparkar, with each taluka comprising 70 (n=70) water samples (250 ml each) collected randomly in December 2021 and May 2021. The samples were aseptically packed into the sterile bottles and transported to the laboratory in ice boxes and then immediately processed or stored at 4°C 24-48 hours in the event of delayed analysis. To isolate Gram-negative bacilli, such as the coliforms, in the sample water collected, the Filtration Assembly Method (FAM) was carried out on each sample [11]. The filtration process was done through the membrane filtration assembly (Millipore, USA) unit of a funnel, vacuum pump, filter flask, and filter membrane (Merck, USA) with pore sizes of 0.45 micrometer (μm). An autoclave was used to sterilize the membrane filtration assembly at 121°C for up to 15-20 minutes. Assembled aseptically Sterile membrane filter unit was assembled. With Filtration Assembly, the water samples (100 ml) were filtered with the assistance of sterile forceps, removing the membrane filter and placing it on Chromatic™ Coli Coliform (Liofilchem, Italy) agar and on MacConkey (Oxoid, UK) agar

plates separately. The Plates were incubated aerobically at 37°C between 24-48 hours. This was done to all the water samples. The plates were then incubated, and the appearance of colored colonies was noted, which validated the findings of the Chromogenic agar of the specific bacteria that were recommended. Moreover, the antimicrobial susceptibility profile was performed based on the principles of Clinical and Laboratory Standards Institute (CLSI 2020) [15] using Kirby-Bauer disc diffusion on Muller-Hinton agar. The antibiotic discs used were Oxoid (UK) which included Amoxicillin plus clavulanic acid (AMC, 30+10 μg) or augmentin (AMC) and Ceftazidime (CAZ, 30 μg) and Cefotaxime (CTX, 30 μg) were both 3rd generation cephalosporin, Aztreonam (ATM, 30 μg) and Ciprofloxacin (CIP, 5 μg). Selection of bacterial colonies was done and bacterial lawn was prepared on Muller Hinton agar plates. After the antibiotic discs had been placed, the plate was incubated at 37°C for 24 hours. The outcome of the antibiotic susceptibility test, as given by the CLSI guidelines, is expressed as sensitive(S) or resistant(R).

RESULTS

Pure bacterial cultures were first isolated from drinking water using chromogenic agar. The filter paper placed onto the chromogenic agar plate surface indicates the growth of specific bacterial colonies. Indications of the appearance of turquoise-blue colonies of *E. coli* and for *Klebsiella* mauve colonies were seen on chromogenic agar (Chromatic™ Coli Coliform) as mentioned by the manufacturer. While some other colorless colonies (Creamy/off-white color) of other bacterial species belong to the family Enterobacteriaceae (Figure 1).



Figure 1: Isolation of Pure Bacterial Culture on Chromogenic Agar
As a result of overnight incubation after FAM, growth was observed. The colonies of lactose-fermenting Coliform group of bacteria appeared as medium to large pink colonies. However, non-lactose fermenting non-Coliform bacteria appeared as colorless colonies that need further identification based on biochemical characterization. From these colonies, pure cultures were obtained by the streak plate method (Figure 2).



Figure 2: Growth on MacConkey Agar and Subculture of Pure Isolates on MacConkey

Out of 70 samples collected from 5 different talukas of the district Thararkar, 47 (70 %) samples were positive for bacterial growth, whereas 23 samples (30%) were found negative. Among 47 positive samples, 28 samples (60 %) were positive for the presence of Coliform, and 19 (40%) samples showed the presence of non-coliform bacteria (Figure 3).




Figure 3: Distribution of Water Samples for Bacterial Contamination (A) Bacterial Distribution of Water Samples (B) Coliform and Non-Coliform Distribution

Other than Coliform bacteria, non-lactose-fermenting, Gram-negative bacteria were also isolated, including *Salmonella* species. Further investigation, biochemical analysis of isolated pure culture results showed that five types of Gram-negative bacilli were isolated, including *E. coli* 18 (40%), *Klebsiella* species 12 (27%), members of coliform and *Pseudomonas* spp. 9 (20%), *Salmonella* species, 3 (6%), and *Proteus* spp. 3 (6%) members of the non-coliform group of bacteria (Table 1).

Table 1: Distribution of Gram-negative Bacilli Isolated

Coliform (n=30)		Non-Coliform (n=15)		
<i>E. coli</i>	<i>Klebsiella</i> spp	<i>Salmonella</i> spp	<i>Pseudomonas</i> spp	<i>Proteus</i> spp
18(40%)	12(27%)	3(6%)	9(20%)	3(6%)

For antibiotics sensitivity pattern, following 24-h incubation, 29(93%) isolates were resistant to Amoxicillin, 21 (68%) were resistant to Cefotaxime and 19 (60%) were resistant to Ceftazidime with 2 (7%), 10 (32%) and 11 (40%) being sensitive, respectively. However, Imipenem,

Ciprofloxacin, and Fosfomycin showed less or no resistance with 93% sensitivity and 7% resistance (Table 2).

Table 2: Sensitivity Pattern of Bacterial Isolates

Sr. No.	Antibiotics	Resistant (R)	Percentage (%)	Sensitive (S)	Percentage (%)
1	AMC	29	93	2	7
2	CAZ	19	60	11	40
3	CTX	21	68	10	32
4	ATM	19	60	11	40
5	CIP	2	7	29	93
6	IPM	2	7	29	93
7	FOS	2	7	29	93

DISCUSSION

Chromatic™ Coli Coliform has chromogenic that is used by the bacteria to produce colored colonies, in which substrate is a chromogen, X-glucuronidase, and through which we can be able to observe different bacterial species. The colonies of *E. coli* were observed as blue in color since the X-glucuronidase cleaved the enzyme chromogen, X-glucuronidase. This medium is specifically applicable in the identification of coliforms and the identification of *E. coli* and *Klebsiella* spp. In our study, 47 samples were positive with bacterial growth of the total 70 samples of water found in the Thar area, and 28 of the 47 positive samples were coliforms. A similar trend was observed in other research, wherein 71 percent of *E. coli* isolated in drinking water out of 155 samples [15]. Sood, N conducted a study on the Prevalence of pathogenic bacteria on fecal coliforms. 50 water samples were sampled; *E. coli* was isolated from these samples 12 [16]. A similar study was conducted locally in Aga Khan university Karachi, Pakistan, and 42 water samples were sampled, in which 62.96% of the samples were found to be *E. coli* [17]. Antibiotic, levofloxacin, has been reported to be most sensitive to the *E. coli* and *Salmonella*, and the most resistant was reported to be against antibiotics clindamycin and amoxicillin [18, 19]. Similar findings have been brought to light in this study, with maximum resistance (93%) against the more popularized antibiotic Augmentin/Amoxicillin, and moreover against third-generation cephalosporin members like cefotaxime and ceftazidime, 68% and 60% respectively. This is a worrying scenario to Thar people who have fewer hospitals and a deficiency of medicine. Our water isolates have a high prevalence of resistance to frontline antibiotics such as amoxicillin, which is also similar to the critical resistance patterns maintained in clinical pathogens. An investigation into Uganda surgical site infections indicated that 90.9 percent of Enterobacteriaceae isolates were multidrug-resistant, and ampicillin resistance was up to 95.1 percent [20]. This analogy of a compromised ecology of Thar and a

compromised health of Uganda reveals a universal dilemma of resistance caused by the improper use of ordinary antibiotics in healthcare and the household sphere. The principal risk to human health is from drinking contaminated water since it is the most crucial parameter to consider. Out of 70 samples, which were tested in 5 different regions of the district Tharparkar, 47 samples (70 percent) tested positive for coliform/bacterial contamination. It has been noted that the total viable count test conducted on microbial analysis and the sample got excessively contaminated with microbial growth of fecal coliforms and *Escherichia coli* [3]. Again, 18 samples in this research demonstrated the existence of *E. coli*, which is the hallmark of the existence of fecal coliform. Further, the health of the water in the area is not good because it has the Coliform group of bacteria, which is threatening, in case of positive results with coliform presence. The infections may be caused by contaminated water and may be fatal in poor areas as a result of poor medical facilities. In this study, three types of bacteria, which are *E. coli* 18 (40%), *Klebsiella* spp. 12 (27%) and *Salmonella* spp. 3 (6%). The bacteria are causative bacteria of water-borne illness, such as Diarrhea, Dysentery, and Typhoid fever, and can lead to urinary tract infections (UTI). In Pakistan, major illnesses because of contaminated drinking water encompass diarrhoea, gastroenteritis because of Coliform contamination, and typhoid because of *Salmonella typhi*, giardiasis because of intestinal worms *Giardia* spp., and viral hepatitis, like hepatitis A caused by the hepatitis A virus. [5, 10, 21]. It has been observed and hypothesized in recent research that the presence of *E. coli* in water is not necessarily due to feces contamination but may survive and propagate in tropical water. Moreover, the Coliform test remains a useful test of microbiological analysis of drinking water. Species-level identification of other non-coliform bacterial species to be used as part of the current study is a useful tool in the identification of the nature of contamination of water. *Salmonella* spp. was also identified in current research, that are harmful enteric pathogen and not part of the coliform group of bacteria.

CONCLUSIONS

This study shows the presence of coliform bacteria in drinking water of Thar district samples and proves that water samples were not safe for drinking, bathing, or washing purposes, as *E. coli* is a major causative agent for Urinary tract infections. *Escherichia coli* determination and its application in the identification of the major animal source of fecal contamination have been used as an indicator of water fecal contamination, such as *Salmonella* and *Giardia*. Use of Chromogenic agar proves to help in fast recovery and identification of *E. coli*.

Authors Contribution

Conceptualization: SJ

Methodology: SJ, BA

Formal analysis: JML, NQ, HN, S

Writing and Drafting: JML

Review and Editing: BK, JML, SJ, NQ, HN, BA, S

All authors approved the final manuscript and take responsibility for the integrity of the work.

Conflicts of Interest

The authors declare no conflict of interest.

Source of Funding

The author received no financial support for the research, authorship and/or publication of this article.

REFERENCES

- [1] Ashraf M, Hasan FU, Fatima B, Ahsan M. Climate-Smart Water Management Technologies in Pakistan. Inclusive-Smart and Resilient Food Systems and Security. Cham: Springer Nature Switzerland. 2024 Sep; 18: (Pp. 379-394). doi: 10.1007/978-3-031-65968-3_17.
- [2] Memon M, Soomro MS, Akhtar MS, Memon KS. Drinking Water Quality Assessment in Southern Sindh (Pakistan). Environmental Monitoring and Assessment. 2011 Jun; 177(1): 39-50. doi: 10.1007/s10661-010-1616-z.
- [3] Khan MH, Nafees M, Muhammad N, Ullah U, Hussain R, Bilal M. Assessment of Drinking Water Sources for Water Quality, Human Health Risks, and Pollution Sources: A Case Study of the District Bajaur, Pakistan. Archives of Environmental Contamination and Toxicology. 2021 Jan; 80(1): 41-54. doi: 10.1007/s00244-020-00801-3.
- [4] Mehmood S, Ahmad A, Ahmed A, Khalid N, Javed T. Open Access Scientific Reports. 2013; 2(2): 1-7. doi: 10.4172/scientificreports.
- [5] Tufail M, Nasir MJ, Rahman AU, Kakakhel SI, Ashraf M, Tariq A. Community Perceptions of Potable Water Quality and Public Health Implications in Flood-Affected Areas of Nowshera District, Pakistan. World Development Perspectives. 2025 Jun; 38: 100679. doi: 10.1016/j.wdp.2025.100679.
- [6] Bhatti NB, Siyal AA, Qureshi AL. Groundwater Quality Assessment Using Water Quality Index: A Case Study of Nagarparkar, Sindh, Pakistan. Sindh University Research Journal-SURJ(Science Series). 2018 Jun; 50(2): 227-34. doi: 10.26692/sujo/2018.06.0040.
- [7] Ali W, Mushtaq N, Javed T, Zhang H, Ali K, Rasool A et al. Vertical Mixing with Return Irrigation Water the Cause of Arsenic Enrichment in Groundwater of District Larkana Sindh, Pakistan. Environmental

Pollution. 2019 Feb; 245: 77-88. doi: 10.1016/j.envpol. 2018.10.10.

[8] Kumar L, Deitch MJ, Tunio IA, Kumar A, Memon SA, Williams L, et al. Assessment of Physicochemical Parameters in Groundwater Quality of Desert Area (Tharparkar) of Pakistan. Case Studies in Chemical and Environmental Engineering. 2022 Dec; 6: 100232. doi: 10.1016/j.cscee.2022.100232.

[9] Brahman KD, Kazi TG, Afzadi HI, Baig JA, Arain SS, Talpur FN et al. Exposure of Children to Arsenic in Drinking Water in the Tharparkar Region of Sindh, Pakistan. Science of the Total Environment. 2016 Feb; 544: 653-60. doi: 10.1016/j.scitotenv.2015.11.152.

[10] Khan AU, Iqbal Q, Khan FA, Ashraf S. Impact of Urbanization on Potable Water Quality: A Bacteriological Case Study of Bannu City. Ground Water. 2013; 15(1): 76.

[11] El Boujnouni H, Balla KN, Belkadi B, Rahouti M. Comparison Between the Recovery Rate of Three Concentration Protocols of Water Samples Intended for Analysis by Molecular Biology: Membrane Filtration, Filtration on Gauze Pad and Centrifugation. Saudi Journal of Biological Sciences. 2022 Mar; 29(3): 1592-7. doi: 10.1016/j.sjbs.2021.11.004.

[12] Ferone M, Gowen A, Fanning S, Scannell AG. Microbial Detection and Identification Methods: Bench Top Assays to Omics Approaches. Comprehensive Reviews in Food Science and Food Safety. 2020 Nov; 19(6): 3106-29. doi: 10.1111/1541-4337.12618.

[13] Rajapaksha P, Elbourne A, Gangadoo S, Brown R, Cozzolino D, Chapman J. A Review of Methods for the Detection of Pathogenic Microorganisms. Analyst. 2019; 144(2): 396-411.

[14] Salam HA, Ashraf M, Iqbal N, Gul N, Memon S. Beneath the Sands: A Comprehensive Study of Groundwater in Tharparkar Region. Pakistan Council of Research in Water Resources. Islamabad. 2023 Jan; 33-46.

[15] Wayne PA. Clinical and Laboratory Standards Institute. Performance Standards for Antimicrobial Susceptibility Testing. 2011; 13(1): 100-121.

[16] Lyimo B, Buza J, Subbiah M, Smith W, Call DR. Comparison of Antibiotic-Resistant *Escherichia coli* obtained from Drinking Water Sources in Northern Tanzania: A Cross-Sectional Study. Biomedical Center Microbiology. 2016 Nov; 16(1): 254. doi: 10.1186/s12866-016-0870-9.

[17] Sood NK, Patel PC, Patel SM, Mandalia AH. A Study of Prevalence of Pathogenic Bacteria, Particularly, Fecal Coliforms and their Antibiotic Resistance Pattern in Environmental Water Samples of a Tertiary-Care Hospital, Ahmedabad. 2016 Jan; 4(12): 1739-1743. doi: 10.5455/ijmusp.2015.10042015359.

[18] Mir N. Isolation of Multi-Drug-Resistant *Escherichia coli* from Drinking Water of Lahore City, Pakistan. Pakistan Journal of Medical and Health Sciences. 2021 Jan. doi: 10.53350/pjmhs211551080.

[19] Alawally NA, Abouserwel A, Ali MA, Ibrahim HK, Muthanna FM. Antibiotic Sensitivity of Common Bacterial Pathogens Against Levofloxacin. Blood. 2022; 3(6): 6. doi: 10.22271/oral.2022.v8.i3a.1572.

[20] Hope D, Ampaire L, Oyet C, Muwangizi E, Twizerimana H, Apecu RO. Antimicrobial Resistance in Pathogenic Aerobic Bacteria Causing Surgical Site Infections in Mbarara Regional Referral Hospital, Southwestern Uganda. Scientific Reports. 2019 Nov; 9(1): 17299. doi: 10.1038/s41598-019-53712-2.

[21] Jabeen S, Naz H, Kumari N, Naz S. Analysis of Physical, Chemical and Biological Aspects of Drinking Water at University of Sindh Jamshoro. International Journal of Economic and Environmental Geology. 2022 Jan; 13(1): 33-6. doi: 10.46660/ijeeq.v13i1.21.